# Instructions – Video Capsule Endoscopy (VCE aka Pill Camera)

# IT IS VERY IMPORTANT TO BE ON TIME DUE TO TESTING GUIDELINES. LATE ARRIVAL MAY RESULT IN YOUR TEST BEING RE-SCHEDULED.

#### ONE WEEK PRIOR-

Discontinue iron pills or multi-vitamins containing iron.

#### DAY BEFORE PROCEDURE-

<u>VERY IMPORTANT - PLEASE DRINK PLENTY OF WATER THROUGHO</u>UT THE DAY.

- 1. Light breakfast/lunch before 11 AM.
- 2. Start clear liquid diet at 11 AM. [ex. water, gatorade, apple juice, soda, tea, plain black coffee No red or purple, No milk products]
- 3. Nothing by mouth after 10 PM.

### **MORNING OF PROCEDURE-**

- NOTHING BY MOUTH
- Do not apply any lotions, creams or powders to your chest, abdomen, or pelvic area. Wear loose clothing [ex. pajamas or workout clothes, no tight clothing.]
- Do not take any medications.

# **DAY OF PROCEDURE-**

8 AM – You will be connected to the monitor, swallow a large pill, and be sent home with the monitor for the rest of the day. You may resume water immediately. <u>VERY IMPORTANT - PLEASE</u> DRINK PLENTY OF WATER THROUGHOUT THE DAY.

10 AM – Resume clear liquid diet (no red, purple, or milk products) 12:15 PM – May take meds.

4:45 PM – Check-In at the front desk. Please do not arrive before 4:45 pm.

Notes: If you hear beeping, please drink a cup of water. If there is continued beeping, please come back to the office. Any questions, please call us at 325-795-2100. In the event we cannot get to the phone, please call surgery center at 325-795-0053.

# **IMPORTANT INFO FOR AFTER YOUR PROCEDURE:**

- 1. No MRI for 2 weeks following the procedure.
- 2. Contact our office or your primary care doctor's office or go to your nearest emergency room if you experience any severe abdominal pain, severe nausea or vomiting.